arXiv:1301.5594v2 [math.NT] 4 Feb 2013

ABELIAN VARIETIES OVER FIELDS OF FINITE CHARACTERISTIC

YURI G. ZARHIN

Abstract. The aim of this paper is to extend our previous results about Galois action on the torsion points of abelian varieties to the case of (finitely generated) fields of characteristic 2.

1. Introduction

Let K be a field, $\bar{K}$ its algebraic closure, $\bar{K}_s \subset \bar{K}$ the separable algebraic closure of K, $\mathrm{Gal}(K) = \mathrm{Gal}(\bar{K}_s/K) = \mathrm{Aut}(\bar{K}/K)$ the absolute Galois group of K.

Let X be an abelian variety over K. Then we write $\mathrm{End}(X)$ for its ring of K-endomorphisms and $\mathrm{End}^0(X)$ for the finite-dimensional semisimple $\mathbf{Q}$-algebra $\mathrm{End}(X) \otimes \mathbf{Q}$. If n is a positive integer that is not divisible by $\mathrm{char}(K)$ then we write X_n for the kernel of multiplication by n in $X(\bar{K})$; it is well known that X_n is free $\mathbf{Z}/n\mathbf{Z}$-module of rank $2\dim(X)$ [8], which is a Galois submodule of $X(\bar{K}_s)$. We write $\bar{\rho}_{n,X}$ for the corresponding (continuous) structure homomorphism

$$\bar{\rho}_{n,X} : \mathrm{Gal}(K) \to \mathrm{Aut}_{\mathbf{Z}/n\mathbf{Z}}(X_n) \cong \mathrm{GL}(2\dim(X), \mathbf{Z}/n\mathbf{Z}).$$

In particular, if $n = \ell$ is a prime different from $\mathrm{char}(K)$ then X_ℓ is a $2\dim(X)$-dimensional $\mathbf{F}_\ell$-vector space provided with

$$\bar{\rho}_{\ell,X} : \mathrm{Gal}(K) \to \mathrm{Aut}_{\mathbf{F}_\ell}(X_\ell) \cong \mathrm{GL}(2\dim(X), \mathbf{F}_\ell).$$

We write

$$\tilde{G}_\ell = \tilde{G}_{\ell,X,K}$$

for the image (subgroup)

$$\bar{\rho}_{\ell,X}(\mathrm{Gal}(K)) \subset \mathrm{Aut}_{\mathbf{F}_\ell}(X_\ell).$$

By definition, $\tilde{G}_{\ell,X,K}$ is a finite subgroup of

$$\mathrm{Aut}_{\mathbf{F}_\ell}(X_\ell) \cong \mathrm{GL}(2\dim(X), \mathbf{F}_\ell).$$

If $K(X_\ell)$ is the field of definition of all points of order ℓ on X then it is a finite Galois extension of K and the corresponding Galois group $\mathrm{Gal}(K(X_\ell)/K)$ is canonically isomorphic to $\tilde{G}_{\ell,X,K}$. If K'/K is a finite Galois extension of fields then $\mathrm{Gal}(K')$ is a normal open subgroup of finite index in $\mathrm{Gal}(K)$ while $X' = X \times_K K'$ is a $\dim(X)$-dimensional abelian variety over K' and the $\mathrm{Gal}(K')$-modules X_ℓ and X'_ℓ are canonically isomorphic. Under this isomorphism, $\tilde{G}_{\ell,X',K'}$ becomes isomorphic to a certain normal subgroup of $\tilde{G}_{\ell,X,K}$.

This work was partially supported by a grant from the Simons Foundation (#246625 to Yuri Zarkhin).

By functoriality, $\mathrm{End}(X)$ acts on X_n. This action gives rise to the embedding of free $\mathbf{Z}/n\mathbf{Z}$-modules

$$\mathrm{End}(X)\otimes\mathbf{Z}/n\mathbf{Z}\hookrightarrow \mathrm{End}_{\mathbf{Z}/n\mathbf{Z}}(X_n);$$

in addition, the image of $\mathrm{End}(X)\otimes\mathbf{Z}/n\mathbf{Z}$ lies in the centralizer $\mathrm{End}_{\mathrm{Gal}(K)}(X_n)$ of $\mathrm{Gal}(K)$ in $\mathrm{End}_{\mathbf{Z}/n\mathbf{Z}}(X_n)$. Further we will identify $\mathrm{End}(X)\otimes\mathbf{Z}/n\mathbf{Z}$ with its image in $\mathrm{End}_{\mathrm{Gal}(K)}(X_n)$ and write

$$\mathrm{End}(X)\otimes\mathbf{Z}/n\mathbf{Z}\subset \mathrm{End}_{\mathrm{Gal}(K)}(X_n).$$

If ℓ is a prime that is different from $\mathrm{char}(K)$ then we write $T_\ell(X)$ for the $\mathbf{Z}_\ell$-Tate module of X and $V_\ell(X)$ for the corresponding $\mathbf{Q}_\ell$-vector space

$$V_\ell(X)=T_\ell(X)\otimes_{\mathbf{Z}_\ell}\mathbf{Q}_\ell$$

provided with the natural continuous Galois action [12]

$$\rho_{\ell,X}:\mathrm{Gal}(K)\to \mathrm{Aut}_{\mathbf{Z}_\ell}(T_\ell(X))\subset \mathrm{Aut}_{\mathbf{Q}_\ell}(V_\ell(X)).$$

Recall [8] that $T_\ell(X)$ is a free $\mathbf{Z}_\ell$-module of rank $2\mathrm{dim}(X)$ and $V_\ell(X)$ is a $\mathbf{Q}_\ell$-vector space of dimension $2\mathrm{dim}(X)$. Notice that there are canonical isomorphisms of $\mathrm{Gal}(K)$-modules

$$X_{\ell^i}=T_\ell(X)/\ell^i T_\ell(X) \tag{0}$$

for all positive integers i. The natural embeddings

$$\mathrm{End}(X)\otimes\mathbf{Z}/\ell^i\mathbf{Z}\hookrightarrow \mathrm{End}_{\mathbf{Z}/\ell^i\mathbf{Z}}(X_{\ell^i})$$

are compatible and give rise to the embeddings of $\mathbf{Z}_\ell$-algebras

$$\mathrm{End}(X)\otimes\mathbf{Z}_\ell\hookrightarrow \mathrm{End}_{\mathbf{Z}_\ell}(T_\ell(X))$$

and $\mathbf{Q}_\ell$-algebras

$$\mathrm{End}(X)\otimes\mathbf{Q}_\ell\hookrightarrow \mathrm{End}_{\mathbf{Q}_\ell}(V_\ell(X)).$$

Again, the images of $\mathrm{End}(X)\otimes\mathbf{Z}_\ell$ in $\mathrm{End}_{\mathbf{Z}_\ell}(T_\ell(X))$ and of $\mathrm{End}(X)\otimes\mathbf{Q}_\ell$ in $\mathrm{End}_{\mathbf{Q}_\ell}(V_\ell(X))$ lie in the centralizers $\mathrm{End}_{\mathrm{Gal}(K)}(T_\ell(X))$ and $\mathrm{End}_{\mathrm{Gal}(K)}(V_\ell(X))$ respectively. We will identify $\mathrm{End}(X)\otimes\mathbf{Z}_\ell$ and $\mathrm{End}(X)\otimes\mathbf{Q}_\ell$ with their respective images and write

$$\mathrm{End}(X)\otimes\mathbf{Z}_\ell\subset \mathrm{End}_{\mathrm{Gal}(K)}(T_\ell(X))\subset \mathrm{End}_{\mathbf{Z}_\ell}(T_\ell(X)),$$

$$\mathrm{End}(X)\otimes\mathbf{Q}_\ell\subset \mathrm{End}_{\mathrm{Gal}(K)}(V_\ell(X))\subset \mathrm{End}_{\mathbf{Q}_\ell}(V_\ell(X)).$$

Similarly, if Y is (may be) another abelian variety over K then we write $\mathrm{Hom}(X,Y)$ for the (free commutative) group of all K-homomorphisms from X to Y. Similarly, there are the natural embeddings

$$\mathrm{Hom}(X,Y)\otimes\mathbf{Z}/n\mathbf{Z}\subset \mathrm{Hom}_{\mathrm{Gal}(K)}(X_n,Y_n)\subset \mathrm{Hom}_{\mathbf{Z}/n\mathbf{Z}}(X_n,Y_n),$$

$$\mathrm{Hom}(X,Y)\otimes\mathbf{Z}/\ell^i\mathbf{Z}\subset \mathrm{Hom}_{\mathrm{Gal}(K)}(X_{\ell^i},Y_{\ell^i})\subset \mathrm{Hom}_{\mathbf{Z}/\ell^i\mathbf{Z}}(X_{\ell^i},Y_{\ell^i}),$$

$$\mathrm{Hom}(X,Y)\otimes\mathbf{Z}_\ell\subset \mathrm{Hom}_{\mathrm{Gal}(K)}(T_\ell(X),T_\ell(Y))\subset \mathrm{Hom}_{\mathbf{Z}_\ell}(T_\ell(X),T_\ell(Y)),$$

$$\mathrm{Hom}(X,Y)\otimes\mathbf{Q}_\ell\subset \mathrm{Hom}_{\mathrm{Gal}(K)}(V_\ell(X),V_\ell(Y))\subset \mathrm{Hom}_{\mathbf{Q}_\ell}(V_\ell(X),V_\ell(Y)).$$

Let

$$\rho_{\ell,X}:\mathrm{Gal}(K)\to \mathrm{Aut}_{\mathbf{Z}_\ell}(T_\ell(X))\subset \mathrm{Aut}_{\mathbf{Q}_\ell}(V_\ell(X))$$

be the corresponding ℓ-adic representation of $\mathrm{Gal}(K)$. The image

$$G_{\ell,X}=\rho_{\ell,X}(\mathrm{Gal}(K))\subset \mathrm{Aut}_{\mathbf{Z}_\ell}(T_\ell(X))\subset \mathrm{Aut}_{\mathbf{Q}_\ell}(V_\ell(X))$$

is a compact ℓ-adic Lie (sub)group [11, 12].

Let d be a positive integer. We write $\mathrm{Isog}(X,K,d)$ for the set of K-isomorphism classes of abelian varieties Y over K that enjoy the following properties:

(i) Y admits a K-polarization of degree d;
(ii) There exists a K-isogeny $Y \to X$, whose degree is prime to $\mathrm{char}(K)$.

For example, if $d=1$ then $\mathrm{Isog}(X,K,1)$ consists of (K-isomorphism classes of) all principally polarized abelian varieties Y over K that admit a K-isogeny, whose degree is prime to $\mathrm{char}(K)$.

We write $\mathrm{Isog}(X,K)$ for the set of K-isomorphism classes of abelian varieties Y over K such that there exists a K-isogeny $B \to A$, whose degree is prime to $\mathrm{char}(K)$. Clearly, $\mathrm{Isog}(X,K)$ coincides with the union of all $\mathrm{Isog}(X,K,d)$'s.

The following statement was proven under an additional assumption that p does not divide d by the author when $p>2$ [17] and by S. Mori when $p=2$ [7, Ch. XII, Cor. 2.4 on p. 244]. (This is a strenghening of Tate' finiteness conjecture for isogenies of abelian varieties [15, 23].)

Theorem 1.1. *Assume that* $p:=\mathrm{char}(K)>0$ *and* K *is finitely generated over the finite prime field* $\mathbf{F}_p$. *Let* d *be a positive integer and* A *be an abelian variety over* K.

Then the set $\mathrm{Isog}(A,K,d)$ *is finite.*

The finiteness of $\mathrm{Isog}(A,K,d)$ combined with results of [16] implies the Tate conjecture on homomorphisms of abelian varieties and the semisimplicity of Tate modules over K (see [17] for $p>2$ and [7, Ch. XII, Th. 2.5 on pp. 244–245]).

Theorem 1.2. *Assume that* $p:=\mathrm{char}(K)>0$ *and* K *is finitely generated over the finite prime field* $\mathbf{F}_p$.

Then for all abelian varieties A *and* B *over* K *and every prime* $\ell \ne \mathrm{char}(K)$ *the Galois module* $V_\ell(A)$ *is semisimple and the natural embedding of* $\mathbf{Z}_\ell$*-modules*

$$\mathrm{Hom}_K(A,B)\otimes \mathbf{Z}_\ell \hookrightarrow \mathrm{Hom}_{\mathrm{Gal}(K)}(T_\ell(A),T_\ell(B)$$

is bijective.

By Lemma 1 of [15, Sect. 1] the second assertion of Theorem 1.2 implies the following statement.

Theorem 1.3. *Assume that* $p:=\mathrm{char}(K)>0$ *and* K *is finitely generated over the finite prime field* $\mathbf{F}_p$.

Then for all abelian varieties A *and* B *over* K *the natural embedding of* $\mathbf{Q}_\ell$*-vector spaces*

$$\mathrm{Hom}_K(A,B)\otimes \mathbf{Q}_\ell \hookrightarrow \mathrm{Hom}_{\mathrm{Gal}(K)}(V_\ell(A),V_\ell(B)$$

is bijective.

1.4. Let K be a field that is finitely generated over the finite prime field $\mathbf{F}_p$ and A an abelian variety of positive dimension over K. Let ℓ be a prime different from p. By Theorem 1.3 (applied to $B=A$) and Theorem 1.2, the $\mathrm{Gal}(K)$-module $V_\ell(A)$ is semisimple and

$$\mathrm{End}_{\mathrm{Gal}(K)}(V_\ell(A))=\mathrm{End}(A)\otimes\mathbf{Q}_\ell=\mathrm{End}^0(A)\otimes_{\mathbf{Q}}\mathbf{Q}_\ell.$$

Since $G_{\ell,A,K}$ is the image of $\mathrm{Gal}(K)\to \mathrm{Aut}_{\mathbf{Z}_\ell}(T_\ell(A))\subset \mathrm{Aut}_{\mathbf{Q}_\ell}(V_\ell(A))$, the $G_{\ell,A,K}$-module $V_\ell(A)$ is semisimple and

$$\mathrm{End}_{G_{\ell,A,K}}(V_\ell(A))=\mathrm{End}(A)\otimes\mathbf{Q}_\ell=\mathrm{End}^0(A)\otimes_{\mathbf{Q}}\mathbf{Q}_\ell.$$

Let $\mathbf{Q}_\ell G_{\ell,A,K}$ be the $\mathbf{Q}_\ell$-subalgebra of $\mathrm{End}_{\mathbf{Q}_\ell}(V_\ell(A))$ spanned by $G_{\ell,A,K}$. It follows from the Jacobson density theorem that $\mathbf{Q}_\ell G_{\ell,A,K}$ coincides with the centralizer of $\mathrm{End}(A)\otimes\mathbf{Q}_\ell$ in $\mathrm{End}_{\mathbf{Q}_\ell}(V_\ell(A))$. It follows easily that if $\mathbf{Z}_\ell G_{\ell,A,K}$ is the $\mathbf{Z}_\ell$-subalgebra of $\mathrm{End}_{\mathbf{Z}_\ell}(T_\ell(A))$ spanned by $G_{\ell,A,K}$ then the centralizer of $\mathrm{End}(A)\otimes\mathbf{Z}_\ell$ in $\mathrm{End}_{\mathbf{Z}_\ell}(T_\ell(A))$ contains $\mathbf{Z}_\ell G_{\ell,A,K}$ as a $\mathbf{Z}_\ell$-submodule of finite index.

2. Main results

The aim of this note is to prove variants of Theorem 1.1 where Tate modules are replaced by Galois modules A_n and B_n. Most of our results were already proven in [18, 28] or stated in [21] under an additional assumption $p>2$. (See also [27] where the case of finite fields is discussed.)

Throughout this section, K is a field that is finitely generated over the finite prime field $\mathbf{F}_p$.

Theorem 2.1. *Let A be an abelian variety of positive dimension over K. Then the set* $\mathrm{Isog}(A,K)$ *is finite.*

Remark 2.2. A weaker version of Theorem 2.1 (where $\mathrm{Isog}(A,K)$ is replaced by its subset that consists of abelian varieties that admit a polarization of degree prime to p) is proven in [18, Th. 6.1] under an additional assumption that $p>2$.

Corollary 2.3. *Let A be an abelian variety of positive dimension over K. There exists a positive integer $r=r(A)$ that is not divisible by p and enjoys the following properties.*

- (i) *If C is an abelian variety over K that admits a K-isogeny $C\to A$, whose degree is not divisible by p then there exists a K-isogeny $\beta: A\to C$ with $\ker(\beta)\subset A_r$.*
- (ii) *If n is a positive integer that is not divisible by p and W is a Galois submodule in A_n then there exists $u\in\mathrm{End}(A)$ such that*
$$rW\subset u(A_n)\subset W.$$
- (iii) *For all but finitely many primes ℓ the Galois module A_ℓ is semisimple.*

Remark 2.4. Corollary 2.3(iii) is proven in [18, Th. 1.1] under an additional assumption $p>2$.

Theorem 2.5. *Let A be an abelian variety over K. Then there exists a positive integer $r_1=r_1(A)$ that enjoys the following properties.*

Let n be a positive integer that is not divisible by p and u_n an endomorphism of the Galois module A_n. If we put $m:=n/(n,r_1)$ then there exists $u\in\mathrm{End}(A)$ such that both u and u_n induce induce the same endomorphism of the Galois module A_m.

If A and B are abelian varieties over K then applying Theorem 2.5 to their product $C=A\times B$, we obtain the following statement.

Theorem 2.6. *Let A and B be abelian varieties over K. Then there exists a positive integer $r_2=r_2(A,B)$ that enjoys the following properties.*

Let n be a positive integer that is not divisible by p and $u_n: A_n\to B_n$ a homomorphism of the Galois modules. If we put $m:=n/(n,r_1)$ then there exists $u\in\mathrm{Hom}(A,B)$ such that both u and u_n induce the same homomorphism of the Galois modules $A_m\to B_m$.

Theorem 2.6 implies readily the following assertion.

Corollary 2.7. *Let A and B be abelian varieties over K. Then for all but finitely many primes ℓ the natural injection*

$$\mathrm{Hom}(A,B)\otimes \mathbf{Z}/\ell\mathbf{Z} \hookrightarrow \mathrm{Hom}_{\mathrm{Gal}(K)}(A_\ell, B_\ell)$$

is bijective.

Remark 2.8. Theorem 2.6 was stated without proof in [21] under an additional condition that $p>2$. Corollary 2.7 was proven in [18, Th. 1.1] under an additional condition that $p>2$.

Theorem 2.9. *Let A be an abelian variety of positive dimension over K. Then for all but finitely many primes ℓ the centralizer of* $\mathrm{End}(A)\otimes\mathbf{Z}_\ell$ *in* $\mathrm{End}_{\mathbf{Z}_\ell}(T_\ell(A))$ *coincides with* $\mathbf{Z}_\ell G_{\ell,A,K}$.

Remark 2.10. When K is a field of characteristic zero that is finitely generated over $\mathbf{Q}$, an analogue of Theorem 2.9 was proven by G. Faltings [3, Sect. 3, Th. 1(c)].

Recall that an old result of Grothendieck [9] asserts that in characteristic p an abelian variety of CM-type is isogenous to an abelian variety that is defined over a finite field. (The converse follows from a theorem of Tate [15].)

Theorem 2.11. *Let A be an abelian variety over K. Suppose that for infinitely many primes ℓ the group $\tilde{G}_{\ell,X,K}$ is commutative. Then A is an abelian variety of CM type over K and therefore is isogenous over $\bar{K}$ to an abelian variety that is defined over a finite field.*

Theorem 2.11 may be strengthened as follows.

Theorem 2.12. *Let A be an abelian variety over K. Suppose that for infinitely many primes ℓ the group $\tilde{G}_{\ell,X,K}$ is ℓ-solvable, i.e., its Jordan–Hölder factors are either ℓ-groups or groups, whose order is not divisible by ℓ. Then there is a finite Galois extension K'/K such that $A\times_K K'$ is an abelian variety of CM type over K' and therefore is isogenous over $\bar{K}$ to an abelian variety that is defined over a finite field.*

Theorem 2.12 combined with the celebrated theorem of Feit-Thompson (about solvability of groups of odd order) implies readily the following statement.

Corollary 2.13. *Let A be an abelian variety of positive dimension over K that is not isogenous over $\bar{K}$ to an abelian variety that is defined over a finite field. Then for all but many primes ℓ the group $\tilde{G}_{\ell,X,K}$ is not solvable and its order is divisible by 2ℓ.*

Remark 2.14. See [24, 26, 25] for a plenty of explicit examples of abelian varieties in characteristic p without CM.

Theorem 2.11 was proven in [19] under an additional assumption that $p>2$. Theorem 2.12 was stated without proof in [21] under an additional assumption that K is global and $p>2$.

Theorem 2.15. *Let $K^{ab}\subset \bar{K}_s$ be the maximal abelian extension of K, whose degree is prime to p. If X ia an abelian variety over K such that* $\mathrm{End}(X\times_K\bar{K})$ *is an order in a direct sum of totally real number fields then the torsion subgroup of $X(K^{ab})$ is finite.*

Remark 2.16. Theorem 2.15 was proven in [19, Th. 1.5] under an additional assumption that $p > 2$.

The paper is organized as follows. In Section 3 we discuss isogenies of abelian varieties and their kernels (viewed as finite Galois modules). One of the goal of our approach is to stress the role of analogues of Tate's finiteness conjecture for isogeny classes of abelian varieties. In Section 4 we prove all the main results except Theorem 2.12, which will be proven in Section 5. Section 6 contains additional references to results that may be extended to characteristic 2 case.

3. Isogenies and finite Galois modules

We write $\mathbf{P}$ for the set of all primes. Let K be a field. Let $P \subset \mathbf{P}$ be a nonempty set of primes that does *not* contain $\mathrm{char}(K)$. If A and B are abelian varieties over K then a K-isogeny $A \to B$ is called a P-isogeny if all prime divisors of its degree are elements of P. For example, if P is a singleton $\{\ell\}$ then a P-isogeny is nothing else but an ℓ-isogeny. We say that A and B are P-isogenous over K if there is a P-isogeny $u : A \to B$ that is defined over K. The property to be P-isogenous is an equivalence relation. Indeed, one has only to check that there exists a P-isogeny $v : B \to A$ that is defined over K. Indeed, thanks to Lagrange theorem, $\ker(u) \subset A_n$ where

$$n := \deg(u) = \#(\ker(u)).$$

It follows that there is a K-isogeny $v : B \to A$ such that the composition $vu : A \to B \to A$ coincides with multiplication by n in A. This implies that

$$n^{2\dim(A)} = \deg(vu) = \deg(v)\deg(u).$$

Since u is a P-isogeny, all prime divisors of n belong to P. This implies that all prime divisors of $\deg(v)$ also belong to P, i.e., v is a P-isogeny and we are done.

Let A^t and B^t be the dual abelian varieties (over K) of A and B respectively and

$$u^t : B^t \to A^t, \ v^t : A^t \to B^t$$

be the K-isogenies that are duals of u and v respectively. Since

$$\deg(u^t) = \deg(u). \deg(v^t) = \deg(v),$$

A^t and B^t are also P-isogenous over K. (Warning: A and A^t are *not* necessarily P-isogenous!) This implies that if A and B are P-isogenous over K then $(A \times A^t)^4$ and $(B \times B^t)^4$ are also P-isogenous over K.

We write $\mathrm{Isog}_P(A, K)$ for the set of isomorphism classes of abelian varieties B over K that are P-isogenous to A over K. We write $\mathrm{Isog}_P(A, K, 1)$ for the subset of $\mathrm{Isog}_P(A, K)$ that consists of all isomorphism classes of B with principal polarization over K. For example, if P is $\mathbf{P} \setminus \{\mathrm{char}(K)\}$ then

$$\mathrm{Isog}_P(A, K) = \mathrm{Isog}(A, K), \ \mathrm{Isog}_P(A, K, 1) = \mathrm{Isog}(A, K, 1).$$

Now Theorem 2.1 becomes an immediate corollary of Theorem 1.1 and the following statement.

Theorem 3.1. *Let A be an abelian variety over a field K. Suppose that the set $\mathrm{Isog}_P((A \times A^t)^4, K, 1)$ is finite. Then the set $\mathrm{Isog}_P(A, K)$ is also finite.*

Proof. (i) Let us fix an abelian variety A be over K. Let B be an abelian variety over K that is P-isogenous to A over K. As we have seen, $(B\times B^t)^4$ is P-isogenous to $(A\times A^t)^4$ over K. Recall [18, 22, 7] (see also [27, Sect. 7] that $(B\times B^t)^4$ admits a principal polarization over K. Since the set $\mathrm{Isog}_P((A\times A^t)^4, K, 1)$ is finite, the set of K-isomorphism classes of all $(B\times B^t)^4$ (with fixed A) is finite. On the other hand, each B is isomorphic to a K-abelian subvariety of $(B\times B^t)^4$ over K. But the set of isomorphism classes of abelian subvarieties of a given abelian variety is finite [6]. This implies that the set of K-isomorphism classes of all B's is finite □

Corollary 3.2. *Let X be an abelian variety of positive dimension over a field K. Suppose that the set $\mathrm{Isog}_P((X\times X^t)^4, K, 1)$ is finite.*

Then there exists a positive integer $r=r(X)$ that is not divisible by $\mathrm{char}(K)$ and enjoys the following properties.

- (i) *If Y is an abelian variety over K that is P-isogenous to X over K then there exist a P-isogeny $\beta: X\to Y$ over K with $\ker(\beta)\subset A_r$.*
- (ii) *If n is a positive integer, all whose prime divisors lie in P and W is a Galois submodule in A_n, then there exists $u\in \mathrm{End}(A)$ such that*

$$rW\subset u(A_n)\subset W.$$

Proof. By Theorem 3.1, there are finitely many K-abelian varieties $Y_1,\dots,Y_d$ that are P-isogenous to X over K and such that every K-abelian variety Y over K that is P-isogenous to X over K is K-isomorphic to one of Y_j. For each C_i pick a P-isogeny $\beta_i: X\to Y_i$ that is defined over K. Clearly, $\ker(\beta_i)\subset X_{m_i}$ where $m_i=\deg(\beta_i)$. Let us put $r=\prod_{i=1}^d m_i$. Clearly, for all Y_i

$$\ker(\beta_i)\subset X_{m_i}\subset X_r.$$

This implies that for every K-abelian variety Y over K that is P-isogenous to A over K there exists a P-isogeny $\beta: X\to Y$ over K, whose kernel lies in X_r. This proves (i), since every prime divisor of r is a prime divisor of one of $m_i=\deg(\beta_i)$ and therefore lies in P.

Proof of (ii) The quotient $Y=X/W$ is an abelian variety over K. The canonical map $\pi: X\to X/W=Y$ is a P-isogeny over K, because $\deg(\pi)=\#(W)$ divides $\#(X_n)=n^{2\dim(X)}$. This implies that Y is P-isogenous to X over K.

The rest of the proof goes literally (with the same notation) as in [27, Sect. 8, pp. 331–332] provided one replaces the reference to [27, Cor. 3.5(i)] by the already proven case (i) of Corollary 3.2. (In [27], $n_X: X\to X$ and $n_Y: Y\to Y$ denote the multiplication by n in X and Y respectively.) □

Theorem 3.3. *Suppose that P is infinite. Let X be an abelian variety of positive dimension over a field K. Suppose that the set $\mathrm{Isog}_P((X\times X^t)^4, K, 1)$ is finite.*

Then for all but finitely many primes ℓ the Galois module X_ℓ enjoys the following properties.

If W is a Galois submodule in X_ℓ then there exists $\tilde u\in \mathrm{End}(X)\otimes \mathbf{Z}/\ell Z$ such that $\tilde u^2=\tilde u$ and $\tilde u(X_n)=W$. In particular, the Galois module X_n splits into a direct sum

$$X_n=\tilde u(X_n)\oplus(1-\tilde u)(X_n)=W\oplus(1-\tilde u)(X_n)$$

of its Galois submodules W and $(1-\tilde u)(X_n)$.

Theorem 3.3 implies immediately the following assertion.

Corollary 3.4. *Suppose that P is infinite. Let X be an abelian variety of positive dimension over a field K. Suppose that the set $\mathrm{Isog}_P((X \times X^t)^4, K, 1)$ is finite.*

Then for all but finitely many primes $\ell \in P$ the Galois module X_ℓ is semisimple.

Proof of Theorem 3.3. It is well known that for all but finitely many primes ℓ the finite-dimensional $\mathbf{F}_\ell$-algebra $\mathrm{End}(X) \otimes \mathbf{Z}/\ell\mathbf{Z}$ is semisimple. (See, e.g., [18, Lemma 3.2].) Let r be as in Corollary 3.2. Now let $\ell \in P$ be a prime that does not divide r and such that $\mathrm{End}(X) \otimes \mathbf{Z}/\ell\mathbf{Z}$ is semisimple. Let W be a Galois submodule in X_ℓ. By Corollary 3.2, there exists $u \in \mathrm{End}(X)$ such that

$$rW \subset u(X_\ell) \subset W.$$

Since ℓ does not divide r, we have $rW = W$ and therefore $u(X_\ell) = W$. Let u_ℓ be the image of u in $\mathrm{End}(X) \otimes \mathbf{Z}/\ell\mathbf{Z}$. Clearly,

$$u_\ell(X_\ell) = u(X_\ell) = W.$$

Let I be the right ideal in semisimple $\mathrm{End}(X) \otimes \mathbf{Z}/\ell\mathbf{Z}$ generated by u_ℓ. The semisimplicity implies that there exists an *idempotent* $\tilde{u}$ that generates I. It follows that

$$W = u_\ell(X_\ell) = \tilde{u}(X_\ell).$$

□

We will need the following lemma [27, Lemma 9.2 on p. 333].

Lemma 3.5. *Let Y be an abelian variety of positive dimension over an arbitrary field K. Then there exists a positive integer $h = h(Y, K)$ that enjoys the following properties.*

If n is a positive integer that is not divisible by $\mathrm{char}(K)$, $u, v \in \mathrm{End}(Y)$ are endomorphisms of Y such that

$$\{\ker(u) \bigcap Y_n\} \subset \{\ker(v) \bigcap Y_n\}$$

then there exists a K-isogeny $w : Y \to Y$ such that

$$hv - wu \in n \cdot \mathrm{End}(Y).$$

In particular, the images of hv and wu in

$$\mathrm{End}(Y) \otimes \mathbf{Z}/n\mathbf{Z} \subset \mathrm{End}_{\mathrm{Gal}(K)}(Y_n) \subset \mathrm{End}_{\mathbf{Z}/n\mathbf{Z}}(Y_n)$$

coincide.

Theorem 3.6. *Let X be an abelian variety of positive dimension over a field K. Suppose that the set $\mathrm{Isog}_P((X \times X^t)^8, K, 1)$ is finite. Then there exists a positive integer $r_1 = r_1(X, K)$ that enjoys the following properties.*

Let n be a positive integer, all whose prime divisors lie in P and $m = n/(n, r_1)$. If $u_n \in \mathrm{End}_{\mathrm{Gal}(K)}(X_n)$ then there exists $u \in \mathrm{End}(X)$ such that the images of u_n and u in $\mathrm{End}_{\mathrm{Gal}(K)}(X_m)$ do coincide.

Proof. Let us put $Y = X \times X$. Then $(Y \times Y^t)^4 = (X \times X^t)^8$. Let $r(Y)$ be as in Corollary 3.2 and $h(Y)$ as in Lemma 3.5. Let us put

$$r_1 = r_1(X, K) = r(Y, K)h(Y, K).$$

Now the proof goes literally as the the proof of Theorem 4.1 in [27, Sect. 10], provided one replaces rthe eferences to Cor. 3.5 and Lemma 9.2 of [27] by references to Cor. 3.2 and Lemma 3.5 respectively. □

Let A and B be abelian varieties over K. Applying Theorem to $X = A \times B$ and using the obvious compatible decompositions

$$\mathrm{End}(X) = \mathrm{End}(A) \oplus \mathrm{End}(B) \oplus \mathrm{Hom}(A, B) \oplus \mathrm{Hom}(B, A),$$

$$\mathrm{End}_{\mathrm{Gal}(K)}(X_n) = \mathrm{End}_{\mathrm{Gal}(K)}(A_n) \oplus \mathrm{End}_{\mathrm{Gal}(K)}(B_n) \oplus \mathrm{Hom}_{\mathrm{Gal}(K)}(A_n, B_n) \oplus \mathrm{Hom}_{\mathrm{Gal}(K)}(B_n, A_n),$$

we obtain the following statement.

Theorem 3.7. *Let A and B be abelian varieties of positive dimension over a field K. Suppose that the set $\mathrm{Isog}_P((A \times B \times A^t \times B^t)^8, K, 1)$ is finite. Then there exists a positive integer $r_2 = r_2(A, B, K) = r_1(A \times B, K)$ that enjoys the following properties.*

Let n be a positive integer, all whose prime divisors lie in P and $m = n/(n, r_1)$. If $u_n \in \mathrm{Hom}_{\mathrm{Gal}(K)}(A_n, B_n)$ then there exists $u \in \mathrm{Hom}(A, B)$ such that the images of u_n and u in $\mathrm{Hom}_{\mathrm{Gal}(K)}(A_m, B_m)$ do coincide.

Corollary 3.8. *Let A and B be abelian varieties of positive dimension over a field K. Suppose that the set $\mathrm{Isog}_P((A \times B \times A^t \times B^t)^8, K, 1)$ is finite. Then for all primes $\ell \in P$ the natural injection*

$$\mathrm{Hom}(A, B) \otimes \mathbf{Z}_\ell \hookrightarrow \mathrm{Hom}_{\mathrm{Gal}(K)}(T_\ell(A), T_\ell(B))$$

is bijective.

Proof of Corollary 3.8. Let $r_2 = r_2(A, B)$ be as in Theorem 3.7. Let ℓ^{i_0} be the exact power of ℓ that divides r_2. Let $v \in \mathrm{Hom}_{\mathrm{Gal}(K)}(T_\ell(A), T_\ell(B))$. For each $i > i_0$, v induces a homomorphism $v_i \in \mathrm{Hom}_{\mathrm{Gal}(K)}(A_{\ell^j}, B_{\ell^j})$. By Theorem 3.7, there exists $u_i \in \mathrm{Hom}(A, B)$ such that the images of u_i and v_i in $\mathrm{Hom}(A_{\ell^{i-i_0}}, B_{\ell^{i-i_0}}$. This means that $u_i - v$ sends $T_\ell(A)$ into $\ell^{i-i_0} T_\ell(B)$. It follows that v coincides with the limit of the sequence $\{u_i\}_{i>i_0}^{\infty}$ in $\mathrm{Hom}_{\mathbf{Z}_\ell}(T_\ell(A), T_\ell(B))$ with respect to ℓ-adic topology. Since $\mathrm{Hom}(A, B) \otimes \mathbf{Z}_\ell$ is a compact and therefore a closed subset of $\mathrm{Hom}_{\mathbf{Z}_\ell}(T_\ell(A), T_\ell(B))$, the limit v also lies in $\mathrm{Hom}(A, B) \otimes \mathbf{Z}_\ell$. $\square$

The following lemma will be proven at the end of this section.

Lemma 3.9. *Let X be an abelian variety of positive dimension over a field K. Suppose that the set $\mathrm{Isog}_P((X \times X^t)^4, K, 1)$ is finite. Let $r = r(X, K)$ be as in Corollary 3.2. Then every $\ell \in P$ enjoys the following properties.*

Let $\mathbf{S}$ be a Galois-invariant **pure** *(free) $\mathbf{Z}_\ell$-submodule in $T_\ell(X)$. Then there exists $u \in \mathrm{End}(X) \otimes \mathbf{Z}_\ell$ such that*

$$r_1 \cdot \mathbf{S} \subset u(T_\ell(X)) \subset \mathbf{S}.$$

Theorem 3.10. *Let X be an abelian variety of positive dimension over a field K. Suppose that the set $\mathrm{Isog}_P((X \times X^t)^8, K, 1)$ is finite. Then every $\ell \in P$ enjoys the following properties.*

If $\mathbf{W}$ is a $\mathrm{Gal}(K)$-invariant $\mathbf{Q}_\ell$-vector subspace in $V_\ell(X)$ then there exists $\tilde{u} \in \mathrm{End}(X) \otimes \mathbf{Q}_\ell$ such that $\tilde{u}^2 = \tilde{u}$ and $\tilde{u}(V_\ell(X)) = \mathbf{W}$. In particular, $V_\ell(X)$ splits into a direct sum

$$V_\ell(X) = \tilde{u}(V_\ell(X)) \oplus (1 - \tilde{u})(V_\ell(X)) = \mathbf{W} \oplus (1 - \tilde{u})(V_\ell(X))$$

of its Galois submodules $\mathbf{W}$ and $(1 - \tilde{u})(V_\ell(X))$ and the $\mathrm{Gal}(K)$-module $V_\ell(X)$ is semisimple.

Proof of Theorem 3.10. Let us put $\mathbf{S} = \mathbf{W} \bigcap T_\ell(X)$. Clearly, $\mathbf{S}$ is a pure free Galois-invariant free $\mathbf{Z}_\ell$-submodule in $T_\ell(X)$ and $\mathbf{W} = \mathbf{Q}_\ell\mathbf{S}$. By Lemma 3.9, there exists $u \in \mathrm{End}(X) \otimes \mathbf{Z}_\ell$ such that

$$r \cdot \mathbf{S} \subset u(T_\ell(X)) \subset \mathbf{S}.$$

It follows that

$$r \cdot \mathbf{W} \subset u(V_\ell(X)) \subset \mathbf{W}.$$

Since $r \cdot \mathbf{W} = \mathbf{W}$, we have $u(V_\ell(X)) = \mathbf{W}$. Notice that

$$\mathrm{End}(X) \otimes \mathbf{Z}_\ell \subset \mathrm{End}(X) \otimes \mathbf{Q}_\ell = \mathrm{End}^0(X) \otimes_{\mathbf{Q}} \mathbf{Q}_\ell$$

and $\mathrm{End}^0(X)$ is a finite-dimensional semisimple $\mathbf{Q}$-algebra. It follows that $\mathrm{End}(X) \otimes \mathbf{Q}_\ell$ is a finite-dimensional semisimple $\mathbf{Q}_\ell$-algebra. Let I the left ideal in semisimple $\mathrm{End}(X) \otimes \mathbf{Q}_\ell$ generated by u; there is an idempotent $\tilde{u}$ that generates I. Clearly.

$$\tilde{u}(V_\ell(X)) = u(V_\ell(X)) = \mathbf{W}.$$

□

Proof of Lemma 3.9. If $\mathbf{S} = \{0\}$ then we just put $u = 0$. If $\mathbf{S} = T_\ell(X)$ then we take as u the identity automorphism 1_X of X.

So, further we assume that $\mathbf{S}$ is a proper free $\mathbf{Z}_\ell$-module in $T_\ell(X)$ of positive rank say, d and let $\{e_1, \dots, e_j \dots e_d\}$ be its basis. Since $\mathbf{S}$ is pure in $T_\ell(X)$, for all positive integers i the natural homomorphism of Galois modules

$$\mathbf{S}_i := \mathbf{S}/\ell^i\mathbf{S} \to T_\ell(X)/\ell^i T_\ell(X) = X_{\ell^i}$$

is an injection of free $\mathbf{Z}/\ell^i\mathbf{Z}$-modules. Further, we will identify $\mathbf{S}_i$ with its image in X_{ℓ^i}. We write $\bar{e}^i_j$ for the image of e_j in $\mathbf{S}_i \subset X_{\ell^i}$; clearly, the d-element set $\{\bar{e}^i_j\}_{j=1}^d$ is a basis of the free $\mathbf{Z}/\ell^i\mathbf{Z}$-module $\mathbf{S}_i$. By Corollary 3.2 applied to $n = \ell^i$ and $W = \mathbf{S}_i$ there exists $u_i \in \mathrm{End}(X)$ such that

$$r\mathbf{S}_i \subset u_i(X_{\ell^i}) \subset \mathbf{S}_i.$$

In particular, $u_i(X_{\ell^i})$ contains $r\bar{e}^i_j$ for all j. It is also clear that for each $z \in T_\ell(X)$

$$u_i(z) \in \mathbf{S} + \ell^i T_\ell(X).$$

For each $\bar{e}^i_j$ pick an element

$$\bar{z}^i_j \in T_\ell(X)\ell^i T_\ell(X) = X_{\ell^i}$$

such that $u_i(\bar{z}^i_j) = r\bar{e}^i_j$. Let us pick $z^i_j \in T_\ell(X)$ such that its image in X_{ℓ^i} coincides with $\bar{z}^i_j$. Clearly, the image of $u_i(z^i_j)$ in $T_\ell(X)/\ell^i T_\ell(X) = X_{\ell^i}$ equals $r\bar{e}^i_j$. Using the compactness of $\mathrm{End}(X) \otimes \mathbf{Z}_\ell$ and $T_\ell(X)$, let us choose an infinite increasing sequence of positive integers $i_1 < i_2 < \dots < i_m < \dots$ such that $\{u_{i_m}\}_{m=1}^\infty$ converges in $\mathrm{End}(X) \otimes \mathbf{Z}_\ell$ to some u and $\{z^j_{i_m}\}_{m=1}^\infty$ converges in $T_\ell(X)$ to some z^j for all j with $1 \le j \le d$. It follows that

$$u(z^j) = \lim u_{i_m}(z^j_{i_m}) = re_j.$$

This implies that $u(T_\ell(X)) \supset r \cdot \mathbf{S}$. On the other hand, for each $z \in T_\ell(X)$

$$u_{i_m}(z) \in \mathbf{S} + \ell^{i_m} T_\ell(X).$$

Since $\{i_m\}_{m=1}^\infty$ is an increasing set of positive integers, $\{\mathbf{S} + \ell^{i_m} T_\ell(X)\}_{m=1}^\infty$ is a decreasing set of compact sets, whose intersection is compact $\mathbf{S}$. It follows that $u(z) = \lim u_{i_m}(z)$ lies in $\mathbf{S}$. □

Lemma 3.11. *Let A be an abelian variety of positive dimension over K. Let ℓ be a prime that is different from* $\mathrm{char}(K)$ *and such that the* $\mathrm{Gal}(K)$*-module A_ℓ is semisimple and*

$$\mathrm{End}_{\mathrm{Gal}(K)}(A_\ell) = \mathrm{End}(A) \otimes \mathbf{Z}/\ell\mathbf{Z}.$$

Then the centralizer of $\mathrm{End}(A) \otimes \mathbf{Z}_\ell$ *in* $\mathrm{End}_{\mathbf{Z}_\ell}(T_\ell(A))$ *coincides with* $\mathbf{Z}_\ell G_{\ell,A,K}$.

Proof of Lemma 3.11. Clearly, $\tilde{G}_{A,\ell,K}$ is the image of $\mathrm{Gal}(K) \twoheadrightarrow G_{\ell,A,K} \to \mathrm{Aut}_{\mathbf{F}_\ell}(A_\ell)$. It follows that the $\tilde{G}_{A,\ell,K}$-module A_ℓ is semisimple and

$$\mathrm{End}_{\tilde{G}_{A,\ell,K}}(A_\ell) = \mathrm{End}(A) \otimes \mathbf{Z}/\ell\mathbf{Z}.$$

By the Jacobson density theorem, $\mathbf{F}_\ell\tilde{G}_{A,\ell,K}$ coincides with the centralizer of $\mathrm{End}(A)\otimes \mathbf{Z}/\ell\mathbf{Z}$ in $\mathrm{End}_{\mathbf{F}_\ell}(A_\ell)$. (Here $\mathbf{F}_\ell\tilde{G}_{A,\ell,K}$ is the $\mathbf{F}_\ell$-subalgebra of $\mathrm{End}_{\mathbf{F}_\ell}(A_\ell)$ spanned by $\tilde{G}_{A,\ell,K}$.)

Let M be the centralizer of $\mathrm{End}(A) \otimes \mathbf{Z}_\ell$ in $\mathrm{End}_{\mathbf{Z}_\ell}(T_\ell(A))$. Clearly, M is a pure $\mathbf{Z}_\ell$-submodule of $\mathrm{End}_{\mathbf{Z}_\ell}(T_\ell(A))$ (i.e., the quotient $\mathrm{End}_{\mathbf{Z}_\ell}(T_\ell(A))/M$ is torsion-free); in addition, M contains $\mathbf{Z}_\ell G_{\ell,A,K}$. We have

$$M/\ell M \subset \mathrm{End}_{\mathbf{Z}_\ell}(T_\ell(A)) \otimes_{\mathbf{Z}_\ell} \mathbf{Z}_\ell/\ell\mathbf{Z}_\ell = \mathrm{End}_{\mathbf{F}_\ell}(A_\ell).$$

Clearly, $M/\ell M$ lies in the centralizer of

$$[\mathrm{End}(A) \otimes \mathbf{Z}_\ell] \otimes_{\mathbf{Z}_\ell} \mathbf{Z}_\ell/\ell\mathbf{Z}_\ell = \mathrm{End}(A) \otimes \mathbf{Z}/\ell\mathbf{Z}$$

in $\mathrm{End}_{\mathbf{F}_\ell}(A_\ell)$. This implies that

$$M/\ell M \subset F_\ell\tilde{G}_{A,\ell,K} \subset \mathrm{End}_{\mathbf{F}_\ell}(A_\ell).$$

On the other hand, the image of $\mathbf{Z}_\ell G_{\ell,A,K}$ in

$$\mathrm{End}_{\mathbf{Z}_\ell}(T_\ell(A)) \otimes_{\mathbf{Z}_\ell} \mathbf{Z}_\ell/\ell\mathbf{Z}_\ell = \mathrm{End}_{\mathbf{F}_\ell}(A_\ell)$$

obviously coincides with $\mathbf{F}_\ell\tilde{G}_{A,\ell,K}$. Since this image lies in $M/\ell M$, we conclude that $M/\ell M = \mathbf{F}_\ell\tilde{G}_{A,\ell,K}$ and $M = \mathbf{Z}_\ell G_{\ell,A,K} + \ell \cdot M$. It follows from Nakayama's Lemma that the $\mathbf{Z}_\ell$-module M coincides with its submodule $\mathbf{Z}_\ell G_{\ell,A,K}$. □

4. Proof of main results

Throughout this section, K is a field that is finitely generated over $\mathbf{F}_p$. Let us put $P = \mathbf{P} \setminus \{p\}$.

Proof of Corollary 2.3 and Theorem 2.5. Corollary 2.3 follows readily from Corollary 3.2 combined with Theorem 1.1. Theorem 2.5 follows readily from Theorem 3.6 combined with Theorem 1.1. □

Proof of Theorems 2.1 and 2.6. One has only to combine Theorem 1.1 with Theorems 3.1 and 3.7 respectively. □

Proof of Theorem 2.9. The assertion follows readily from Lemma 3.11 combined with Corollary 2.3 and Theorem 2.5. □

Proof of Theorem 2.15. The proof of [19, Theorem 1.5] works literally provided one replaces the reference to [17] (in [19, Sect. 1.2]) by references to Theorem 1.2 and the reference to [18, Theorem 1.1.1] in [19, Sect. 2.4.1]) by references to Corollaries 2.3 and Corollary 2.7 respectively. □

Proof of Theorem 2.11. The proof of [19, Theorem 4.7.4] works literally provided one replaces the reference to [18, Theorem 1.1.1] by references to Corollaries 2.3 and 2.7. □

5. TORSION AND RAMIFICATION IN SOLVABLE EXTENSIONS

5.1. Let K be an arbitrary field and $\mathcal{O} \subset K$ a discrete valuation ring, whose field of fractions coincides with K. We write $\mathfrak{p}$ for the maximal ideal of $\mathcal{O}$ and p for the characteristic of the residue field $\mathcal{O}/p$. Let L/K be a finite Galois field extension with Galois group $\mathrm{Gal}(L/K)$ and $\mathcal{O}_L$ the integral closure of $\mathcal{O}$ in L. The following assertion is well known (see, e.g., [29, Ch. 5, Sect. 7–10], [10, Ch. 1, Sect. 7]).

(i) $\mathcal{O}_L$ is a principal ideal domain, the set of its maximal ideals is finite and consists of (say) g maximal ideals $\mathfrak{q}_1, \dots, \mathfrak{q}_g$ such that

$$\mathfrak{p}\mathcal{O} = \left(\prod_{i=1}^{t} \mathfrak{q}_i\right)^{e(L/K)}$$

where $e(L/K)$ is the (weak) ramification index at $\mathfrak{p}$. The degree $f(L/K) = [\mathcal{O}_L/\mathfrak{q}_i : \mathcal{O}/p]$ of the field extension $(\mathcal{O}_L/\mathfrak{q}_i)/(\mathcal{O}/p)$ equals the product $f_0 p^s$ where f_0 is the degree of the separable closure of $\mathcal{O}/p$ in $\mathcal{O}_L/\mathfrak{q}_i$ and s is a nonnegative integer that vanishes if and only if $\mathcal{O}_L/\mathfrak{q}_i$ is separable over $\mathcal{O}/p$. The integers f_0 and s do not depend on i. The product

$$e(L/K) \cdot f_0 p^s g = e(L/K) \cdot f \cdot g = [L : K] = \#(\mathrm{Gal}(L/K))$$

In particular, $e(L/K) \cdot p^s$ divides $[L : K]$. The field extension L/K is tamely ramified at p if and only if $\#(I(\mathfrak{q}_i^0))$ is not divisible by p, i.e., $e(L/K)$ is not divisible by p and $\mathcal{O}_L/\mathfrak{q}_i$ is separable over $\mathcal{O}/p$.

(ii) The Galois group $\mathrm{Gal}(L/K)$ acts transitively on the set $\{\mathfrak{q}_i \mid 1 \le i \le g\}$. The corresponding inertia subgroups $I(\mathfrak{q}_i) \subset \mathrm{Gal}(L/K)$ are conjugate subgroups of order $e(L/K)p^s$ in $\mathrm{Gal}(L/K)$. (See [29, Ch. 5, Sect. 10, Th. 24 and its proof].)

(iii) Let L_0/K be a Galois subextension of L/K, i.e., L_0/K is a Galois field extension and $L_0 \subset L$. Then $\mathfrak{q}_i^0 = \mathfrak{q}_i \bigcap \mathcal{O}_{L_0}$ is a maximal ideal in $\mathcal{O}_{L_0}$, which lies above $\mathfrak{p}$. The image of $I(q_i)$ under the surjection $\mathrm{Gal}(L/K) \twoheadrightarrow \mathrm{Gal}(L_0/K)$ coincides with the inertia group $I(\mathfrak{q}_i^0)$ [10, Ch. 1, Sect. 7, Prop. 22(b)]; in particular, $\#(I(\mathfrak{q}_i^0))$ divides $\#(I(q_i))$. On the other hand, $\#(I(\mathfrak{q}_i^0))$ divides $\#(\mathrm{Gal}(L_0/K)) = [L_0 : K]$. This implies that if $[L_0 : K]$ and $\#(I(q_i))$ are relatively prime then $\#(I(\mathfrak{q}_i^0)) = 1$, i.e., L_0/K is *unramified* at $\mathfrak{p}$.

Let Y be an abelian variety of positive dimension over K. Let $n \ge 3$ be an integer that is not divisible by p. Assume that $Y_n \subset Y(K)$, i.e., all points of order n on X are defined over K.

Let $\mathcal{Y} \to \mathrm{Spec}(\mathcal{O})$ be the Néron model of Y [1]; it is a smooth group scheme, whose generic fiber coincides with Y. Since $Y_n \subset Y(K)$, the Raynaud criterion [5, Prop. 4.7] tells us that Y has *semistable* reduction at $\mathfrak{p}$, i.e., $\mathcal{Y}$ is a semiabelian group scheme.

Let ℓ be a prime different from p. For all positive integers j the field $K(Y_{m_j})$ with $m_j = \ell^j$ is a finite Galois extension. The following assertion was inspired by [7, Ch. XII, Sect. 2.0 on p. 242].

Lemma 5.2. *Let $n \geq 3$ be an integer that is not divisible by p. Assume that $Y_n \subset Y(K)$. Let $L = K(Y_\ell)$ and $\mathfrak{q}$ be a maximal ideal in $\mathcal{O}_L$, which lies above $\mathfrak{p}$. Then:*

- (i) *The inertia group $I(\mathfrak{q})$ is a finite commutative ℓ-group. In particular, L/K is tamely ramified at $\mathfrak{p}$.*
- (ii) *Let L_0/K be a Galois subextension of L/K. If $[L_0 : K]$ is not divisible by ℓ then L_0/K is unramified at $\mathfrak{p}$.*

Proof of Lemma 5.2. The assertion (ii) follows readily from (i). So, let us prove (i).

For all positive integers j let us put $L_j = K(Y_{m_j})$ and $\mathcal{O}_j = \mathcal{O}_{L_j}$. We have $L_1 = L, \mathcal{O}_1 = \mathcal{O}_L$. We have a tower of Galois extensions

$$K \subset L_1 \subset L_2 \subset \dots L_j \subset \dots.$$

We write L_∞ for the union $\bigcup_{i=1}^{\infty} L_i$. For each j pick a maximal ideal $\mathfrak{q}^{(j)}$ in $\mathcal{O}_j$ in such a way that $\mathfrak{q}^{(1)} = \mathfrak{q}$ and $\mathfrak{q}^{(j+1)}$ lies above $\mathfrak{q}^{(j)}$. (Such a choice is possible, because the projective limit of nonempty finite sets is nonempty.) Then the Galois group $\mathrm{Gal}(L_\infty/K)$ is the projective limit of finite groups $\mathrm{Gal}(L_j/K)$'s. It is also clear that

$$\mathrm{Gal}(L_j/K) = \bar{\rho}_{m_j,Y}(\mathrm{Gal}(Y)) \subset \mathrm{Aut}_{\mathbf{Z}/m_j\mathbf{Z}}(Y_{m_j}),$$

$$\mathrm{Gal}(L_\infty/K) = \rho_{\ell,Y}(\mathrm{Gal}(K)) \subset \mathrm{Aut}_{\mathbf{Z}_\ell}(T_\ell(Y)) \subset \mathrm{Aut}_{\mathbf{Q}_\ell}(V_\ell(Y)).$$

Recall that the natural homomorphisms $I(\mathfrak{q}^{(j+1)}) \to I(\mathfrak{q}^{(j)})$ are surjective for all j. Let I_∞ be the projective limit of the corresponding inertia subgroups $I(\mathfrak{q}^{(j)})$; clearly, $I\infty$ is a compact subgroup of $\mathrm{Gal}(L_\infty/K)$ and for each j the natural group homomorphism $I_\infty \to I(\mathfrak{q}^{(j)})$ is surjective, because the projective limit of nonempty finite sets is also nonempty. Therefore one may view I_∞ as a compact subgroup of

$$\mathrm{Aut}_{\mathbf{Z}_\ell}(T_\ell(Y)) \subset \mathrm{Aut}_{\mathbf{Q}_\ell}(V_\ell(Y)).$$

Since Y has semistable reduction at $\mathfrak{q}$, there exists a $\mathbf{Q}_\ell$-vector subspace $\mathbf{W} \subset V_\ell(Y)$ such that I_∞ acts trivially on $\mathbf{W}$ and $V_\ell(Y)/\mathbf{W}$ [5, Prop. 3.5]. It gives us an injective continuous homomorphism of topological groups

$$I_\infty \to \mathrm{Hom}_{\mathbf{Q}_\ell}(V_\ell(Y)/\mathbf{W}), W),$$

$$\sigma \mapsto \{v + W \mapsto \sigma(v) - v\} \ \forall\ \sigma \in I_\infty,\ v \in V_\ell(Y).$$

This implies that there is a continuous isomorphism between I_∞ and its image, which is a compact subgroup of $\mathrm{Hom}_{\mathbf{Q}_\ell}(V_\ell(Y)/\mathbf{W}, \mathbf{W})$. Since the latter is a finite-dimensional $\mathbf{Q}_\ell$-vector space, its every compact subgroup is a commutative pro-ℓ-group (that is isomorphic either to a direct sum of several copies of $\mathbf{Z}_\ell$ or to zero). It follows that I_∞ is also a commutative pro-ℓ-group. Since there is a surjective continuous homomorphism

$$I_\infty \to I(\mathfrak{q}^{(j)}),$$

every $I(\mathfrak{q}^{(j)})$ is a finite commutative ℓ-group. This ends the proof, since $\mathfrak{q} = \mathfrak{q}^{(1)}$ and therefore $I(\mathfrak{q}) = I(\mathfrak{q}^{(1)})$. □

Proof of Theorem 2.12. If K is finite then there is nothing to prove. So, let us assume that K is infinite. Let $d \geq 1$ be the transcendence degree of K over $\mathbf{F}_p$. Let us pick a positive integer $n \geq 3$ that is not divisible by p. Replacing K by $K(X_n)$ and X by $X \times_K K(X_n)$, we may assume that $X_n \subset X(K)$.

Let P be an *infinite* set of primes $\ell \ne p$ such that $\tilde{G}_{\ell,X,K}$ is ℓ-solvable. By deleting finitely many primes from P, we may and will assume that the $\mathrm{Gal}(K)$-module X_{ℓ} is semisimple for all $\ell \in P$. Since $\tilde{G}_{\ell,X,K}$ is the image of $\mathrm{Gal}(K)$ in $\mathrm{Aut}_{\mathbf{F}_{\ell}}(X_{\ell}) \cong \mathrm{GL}(2g, \mathbf{F}_{\ell})$, the $\tilde{G}_{\ell,X,K}$-module X_{ℓ} is semisimple for all $\ell \in P$.

Let $\mathbf{C}$ be the field of complex numbers. Let us put $g = \dim(X)$. Recall that X_{ℓ} is a $2g$-dimensional $\mathbf{F}_{\ell}$-vector space. Let G be a finite ℓ-solvable subgroup of $\mathrm{Aut}_{\mathbf{F}_{\ell}}(X_{\ell})$ such that the natural faithful representation of G in X_{ℓ} is completely reducible. Let us split the semisimple $\mathbf{F}_{\ell}[G]$-module X'_{ℓ} into a direct sum

$$X_{\ell} = \oplus_{i=1}^{m} W_i$$

of simple $\mathbf{F}_{\ell}[G]$-modules W_i. If $d_i = \dim_{\mathbf{F}_{\ell}}(W_i)$ then $2g == \sum_{i=1}^{m} d_i$. By a theorem of Fong-Swan [13, Sect. 16.3, Th. 38], each W_i lifts to characteristic zero; in particular, there is a group homomorphism $\rho_i : G \to \mathrm{GL}(d_i, \mathbf{C})$, whose kernel lies in the kernel of $G \to \mathrm{Aut}_{\mathbf{F}_{\ell}}(W_i)$ (for all i). Clearly, the product-homomorphism

$$\rho = \prod_{i=1}^{d} \rho_i : G \to \prod_{i=1}^{m} \mathrm{GL}(d_i, \mathbf{C}) \subset \mathrm{GL}(2g, \mathbf{C})$$

is an injective group homomorphism and therefore G is isomorphic to a finite subgroup of $\mathrm{GL}(2g, \mathbf{C})$. By a theorem of Jordan [2, Th. Th. 36.13], there is a positive integer $N = N(2g)$ that depends only on $2g$ and such that G contains a normal abelian subgroup of index dividing N. By deleting from P all prime divisors of N, we may and will assume that ℓ does *not* divide N for each $\ell \in P$.

Let us apply this observation to $G = \tilde{G}_{\ell,X,K}$. We obtain that for all $\ell \in P$ the group $\tilde{G}_{\ell,X,K}$ contains an abelian normal subgroup H_{ℓ} of index dividing N. For each $\ell \in P$ let us consider the corresponding subfield of H_{ℓ}-invariants

$$K_{\ell} := (K(X_{\ell}))^{H_{\ell}} \subset K(X_{\ell}).$$

Clearly, K_{ℓ}/K is a Galois extension of degree dividing N while $\mathrm{Gal}(K(X_{\ell})/K_{\ell})$ coincides with commutative (sub)group H_{ℓ}. Since $K_{\ell} \subset K(X_{\ell})$ and $[K_{\ell} : K] = N$ is *not* divisible by ℓ, Lemma 5.2 tells us that the Galois extension K_{ℓ}/K is unramified with respect to every discrete valuation of K. (Since $\mathrm{char}(K) = p$, its every residual characteristic is also p.)

Let k be the (finite) algebraic closure of $\mathbf{F}_p$ in K and let S be an absolutely irreducible normal d-dimensional projective variety over k, whose field of rational functions $k(S)$ coincides with K. Let $V \subset S$ be a *smooth* open dense subset such that the codimension of $S \setminus V$ in S is, at least, 2. Let V_{ℓ} be the normalization of V in K_{ℓ}/K. Now Zariski-Nagata purity theorem tells us that the regular map $V_{\ell} \to V$ is an étale Galois cover; clearly, its degree equals $[K_{\ell} : K]$ and therefore divides N. Let $\pi_1(V)$ be the fundamental group of V that classifies étale covers of V ([4], [7, Ch. XII, Sect. 1, pp. 241–242]. This group is a (natural) topologically finitely generated (topological) quotient of $\mathrm{Gal}(K)$ [7, Ch. XII, Sect. 1, p. 242]and the natural surjection

$$\mathrm{Gal}(K) \twoheadrightarrow \mathrm{Gal}(K_{\ell}/K)$$

factors through $\pi_1(V)$, i.e., it is the composition of the canonical $\mathrm{Gal}(K) \twoheadrightarrow \pi_1(V)$ and a certain continuous surjective homomorphism

$$\gamma_{\ell} : \pi_1(V) \twoheadrightarrow \mathrm{Gal}(K_{\ell}/K),$$

whose kernel is an open normal subgroup of index dividing N.

Since $\pi_1(V)$ is topologically finitely generated, it contains only finitely many open normal subgroups of index dividing N (because it admits only finitely many continuous homomorphisms to any finite group of order dividing N). If Γ is the intersection of all such subgroups then it is an open normal subgroup in $\pi_1(V)$ that lies in the kernel of every γ_ℓ for all $\ell \in P$; in particular, it has finite index. The preimage Δ of Γ is an open normal subgroup of finite index in $\mathrm{Gal}(K)$ and the corresponding subfield of Δ-invariants $E := \bar{K}_s^{\Delta}$ is a finite Galois extension of K that contains K_ℓ for all $\ell \in P$. This implies that for all $\ell \in P$ the compositum $EK(X_\ell)$ is abelian over E , because $K(X_\ell)$ is abelian over K_ℓ. But $EK(X_\ell) = E(X^E{}_\ell)$ where

$$X^E = X \times_K E$$

is an abelian variety over E. Applying Theorem 2.11 to X^E and E (instead of X and K), we conclude that X^E is an abelian variety of CM type and isogenous over $\bar{E} = \bar{K}$ to an abelian variety that is defined over a finite field. The same is true for X, since $X^E = X \times_K E$.

□

6. Concluding Remarks

Theorem 1.2 and Corollaries 2.3(iii) and Corollary 2.7 imply readily that the following results remain true for all prime characteristics p, including $p = 2$.

- Assertions (Sect. 1.3 and 4.4), Corollaries 1–6, Theorem 4.1, and Remark 1 of [19] remain true over any field E that is finitely generated over a finite field of arbitrary characteristic, including 2. Corollary 7 of [19] remains true for any field F of arbitrary prime characteristic, including 2.
- Theorem 1.1(ii) of [14].
- Theorems 1.1, 1.4, 1.6, 1.9, 2.1 of [28].

Acknowledgements. I am grateful to Alexey Parshin, Chad Schoen and Doug Ulmer for their interest in this paper.

References

[1] S. Bosch, W. Lütkebohmert, M. Raynaud, Néron models (Springer-Verlag, Berlin Heidelberg New York, 1990).

[2] Ch. W. Curtis, I. Reiner, Representation theory of finite groups and associative algebras. Interscience Publishers, New York London, 1962.

[3] G. Faltings, ‘Complements to Mordell’, Chapter VI in: G. Faltings, G. Wustholz et al., Rational points. Aspects of Mathematics, E6. Friedr. Vieweg & Sohn, Braunschweig, 1984.

[4] A. Grothendieck et al. , Revêtement étales et groupe fondamental (SGA 1). Lecture Notes in Math. **224**, Springer Verlag, Berlin-Heidelberg-New York, 1971.

[5] A. Grothendieck, *Modèles de Néron et monodromie.* Expose IX dans SGA7 I, Lecture Notes in Math. **288**, Springer Verlag, Berlin-Heidelberg-New York, 1972, pp. 313-523.

[6] H. W. Lenstra, Jr, F. Oort, Yu. G. Zarhin, *Abelian subvarieties.* J. Algebra, **180** (1996), 513–516.

[7] L. Moret-Bailly, Pinceaux de variétés abéliennes, Astérisque, vol. 129 (1985).

[8] D. Mumford, Abelian varieties, Second edition, Oxford University Press, London, 1974.

[9] F. Oort, *The isogeny class of a CM-type abelian variety is defined over a finite extension of the prime field.* J. Pure Applied Algebra **3** (1973), 399–408.

[10] J.-P. Serre, Corps Locaux, Troisieme edition corrigée. Hermann, Paris 1968.

[11] J.-P. Serre, *Sur les groupes des congruence des variétés abéliennes.* Izv. Akad. Nauk SSSR ser. matem. **28**:1 (1964), 3-18; Œuvres **II**, pp. 230–245. Springer-Verlag, Berlin, 1986.

[12] J-P. Serre, Abelian ℓ-adic representations and elliptic curves. Second edition. Addison-Wesley, New York, 1989.

[13] J-P. Serre, Représentations linéares des groupes finis. Troisieme edition corrigée. Hermann, Paris, 1978.

[14] A.N. Skorobogatov, Yu. G. Zarhin, *A finiteness theorem for Brauer groups of abelian varieties and K3 surfaces*. J. Algebraic Geometry **17** (2008), no. 3, 481–502.

[15] J. Tate, Endomorphisms of Abelian varieties over finite fields, Invent. Math. **2** (1966) 134–144.

[16] Yu. G. Zarhin, Endomorphisms of Abelian varieties over fields of finite characteristic,. Izv. Akad. Nauk SSSR ser. matem. **39**(1975) 272–277; Math. USSR Izv. **9** (1975) 255–260.

[17] Yu. G. Zarhin, Abelian varieties in characteristic P, Mat. Zametki **19** (1976) 393–400; Mathematical Notes **19** (1976) 240–244.

[18] Yu. G. Zarhin, Endomorphisms of Abelian varieties and points of finite order in characteristic P, Mat. Zametki **21** (1977) 737–744; Mathematical Notes **21** (1978) 415–419.

[19] Yu. G. Zarhin, Torsion of abelian varieties in fininite characteristic. Mat. Zametki **22** (1977), 1-11; Mathematical Notes **22** (1978), 493 - 498.

[20] Yu. G. Zarhin, *Abelian varieties, ℓ-adic representations and Lie algebras. Rank independence on ℓ*. Invent. Math. **55** (1979), 165–176.

[21] Yu. G. Zarhin, *Homomorphisms of Abelian varieties and points of finite order over fields of finite characteristic* (in Russian), pp. 146-147. In: Problems in Group Theory and Homological Algebra (A. L. Onishchik, editor), Yaroslavl Gos. Univ., Yaroslavl, 1981; MR0709632 (84m:14051).

[22] Yu. G. Zarhin, A finiteness theorem for unpolarized Abelian varieties over number fields with prescribed places of bad reduction, Invent. Math. **79** (1985) 309–321.

[23] Yu. G. Zarhin, A. N. Parshin, *Finiteness problems in Diophantine geometry*. Amer. Math. Soc. Transl. (2) **143** (1989) 35–102; arXiv 0912.4325 [math.NT] .

[24] Yu. G. Zarhin, *Hyperelliptic Jacobians without complex multiplication in positive characteristic*. Math. Research Letters **8** (2001), 429–435.

[25] Yu. G. Zarhin, *Endomorphism rings of certain jacobians in finite characteristic*. Matem. Sbornik **193** (2002), issue 8, 39–48; Sbornik Math., 2002, **193** (8), 1139-1149.

[26] Yu. G. Zarhin, *Non-supersingular hyperelliptic jacobians*. Bull. Soc. Math. France **132** (2004), 617–634.

[27] Yu. G. Zarhin, *Homomorphisms of abelian varieties over finite fields*, pp. 315–343. In: Higher-dimensional geometry over finite fields, (D. Kaledin, Yu. Tschinkel, eds.), IOS Press, Amsterdam, 2008.

[28] Yu. G. Zarhin, *Homomorphisms of abelian varieties over geometric fields of finite characteristic*. J. Inst. Math. Jussieu. Available on CJO 2012 doi:10.1017/S1474748012000655 ; arXiv:1012.5083 [math.NT].

[29] O. Zariski, P. Samuel, Commutative Algebra, vol. I. Van Nostrand, Princeton, NJ 1956.

Department of Mathematics, Pennsylvania State University, University Park, PA 16802, USA

E-mail address: `zarhin@math.psu.edu`